


- $f(4) =$ $f(-3) =$ $f(0) =$ $f(5) =$
 $g(0) =$ $g(x) = 3, x \approx$ $f(x) = 0, x =$ $g(g(1)) \approx$
 $f(g(1)) \approx$ $g(f(5)) \approx$ $f(g(-5)) =$ $f(f(f(\dots f(5)))) =$

f(x)		g(x)		f(4) =	g(3) =	f(0) =
x	y	x	y	f(g(3)) =	g(f(5)) =	f(f(1)) =
1	5	0	3	f(g(5)) =	g(g(4)) =	
2	3	1	1	g(g(...(g(g(1)))) =		
3	2	2	5			
4	1	3	4			
5	4	4	6			
6	4	5	0			



$f(4) = 0$

$f(-3) = \text{DNE}$

$f(0) = 0$

$f(5) = 5$

$g(0) = 1$

$g(x) = 3, x \approx -6.5,$

$f(x) = 0, x = -2, 0, 4 \quad g(g(1)) \approx 2.8$

$f(g(1)) \approx -2.2$

$g(f(5)) \approx -3.8$

$f(g(-5)) = \text{DNE}$

$f(f(f(\dots f(5)))) = 5$

f(x)		g(x)		$f(4) = 1$ $g(3) = 4$ $f(0) = \text{DNE}$ $f(g(3)) = 1$ $g(f(5)) = 6$ $f(f(1)) = 4$ $f(g(5)) = \text{DNE}$ $g(g(4)) = \text{DNE}$ $g(g(\dots(g(g(1)))) = 1$		
x	y	x	y			
1	5	0	3			
2	3	1	1			
3	2	2	5			
4	1	3	4			
5	4	4	6			
6	4	5	0			